

Safety sealed. Do not use if tamper resistant seal around the bottle is broken or missing.

# EMESYL® NAUSEA RELIEF SPRAY

Take at the first signs of nausea symptoms

No known drug interactions

Non-drowsy

No jitters

Non-habit forming

Homepathic

Store at room temperature 15°c - 29°c (59°f - 84°f)

#### **Question? Comment?** Call 1 800 687 1498 or

visit us on the web at www.emesyl.com

Manufactured for Itonis, Incorporated 22951 Mill Creek Dr, Ste A Laguna Hills, CA 92653



**NAUSEA RELIEF SPRAY** 

RELIEF OF NAUSEA AND **NAUSEA** LIKE **SYMPTOMS** 

0.50 FL OZ (15mL) WWW.EMESYL.COM

**BPA-FREE** 



# Drug Facts

## Active Ingredients

Ipecacuanha 3C. Tabacum 3C...... Zingiber officinale 1C.

Nausea symptom relie

Purpose

#### Uses

■ For the symptomatic relief of nausea, and nausea like symptoms.

#### Warnings

### For nasal use only

Ask a doctor before use if you have
■ ear, nose or throat sensitivity ■ susceptibility to nose bleeds

#### When using this product

- Avoid contact with eyes. In case of accidental contact with eyes, flush with water and immediately seek professional help.
   The use of this container by more than one person may spread infection.
- Stop use and ask a doctor if symptoms persist.

If pregnant or breast feeding, ask a health professional before use. Keep out of reach of children. If swallowed, get medical help or contact a

#### Directions

- Pump one or two times into each nostril. Use once every 4 hours as needed.
- Adults and children 6 years of age and older (with adult supervision):
   Remove cap & safety clip (also see illustration on side carton & insert).
   Hold with thumb at bottom of bottle and nozzle between your fingers.
- Before using the first time, prime pump by depressing several times.
   Place tip of nozzle just past nasal opening (approximately 1/8").
   While inside nasal opening, slightly angle nozzle outward.

- Write insue reason opening, signing angle nozze outward.
   Pump one to two times into each nostril.
   After application, press lightly on outside of each nostril for about 5 seconds.
   Wait at least 30 seconds before blowing nose.
   Use once every 4 hours.
- Children under 6 years of age: Consult a doctor before use.

Inactive Ingredients benzalkonium chloride, glycerin, hydroxyethylcellulose, methyl paraben, purified water



8.35"